



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 1/21, 5/10, 15/09, 15/12, 15/63, 15/70, C07K 14/475</b>	<b>A1</b>	<b>(11) International Publication Number: WO 96/39485</b> <b>(43) International Publication Date: 12 December 1996 (12.12.96)</b>
<b>(21) International Application Number:</b> PCT/US95/06731 <b>(22) International Filing Date:</b> 5 June 1995 (05.06.95)  <b>(71) Applicant (for all designated States except US):</b> HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850-3338 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> KUNSCH, Charles, A. [US/US]; 15 Amarillo Court, Germantown, MD 20874 (US). ROSEN, Craig, A. [US/US]; 22400 Rolling Hill Road, Laytonsville, MD 20882 (US).  <b>(74) Agents:</b> SQUIRE, William; Carella, Byrne, Bain, Gilfillan, Cecchi, Stewart & Olstein, 6 Becker Farm Road, Roseland, NJ 07068 (US) et al.		<b>(81) Designated States:</b> AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, ARIPO patent (KE, MW, SD, SZ, UG), European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>With an indication in relation to a deposited microorganism furnished under Rule 13bis separately from the description.</i> <i>Date of receipt by the International Bureau:</i> 4 November 1996 (04.11.96)
<b>(54) Title:</b> HUMAN HEPATOMA-DERIVED GROWTH FACTOR-2  <b>(57) Abstract</b>  A human hepatoma-derived growth factor polypeptide and DNA (RNA) encoding such polypeptide and a procedure for producing such polypeptide by recombinant techniques is disclosed. Also disclosed are methods for utilizing such polypeptide for the stimulation of tissue repair and tissue growth. Antagonists against such polypeptides and their use as a therapeutic to retard tumor growth and scarring is also disclosed. Diagnostic methods for detecting mutations in the coding sequence and alterations in the concentration of the polypeptides in a sample derived from a host are also disclosed.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

## HUMAN HEPATOMA-DERIVED GROWTH FACTOR-2

This invention relates to newly identified polynucleotides, polypeptides encoded by such polynucleotides, the use of such polynucleotides and polypeptides, as well as the production of such polynucleotides and polypeptides. More particularly, the polypeptide of the present invention has been putatively identified as a hepatoma-derived growth factor, sometimes hereinafter referred to "HDGF-2". The invention also relates to inhibiting the action of such polypeptides.

Cell growth is regulated by various growth factors and cytokines, which bind to specific membrane receptors to trigger a cascade of intracellular biochemical signals to the activation of transcription factors, resulting in the activation and repression of various subsets of genes (Aaronson, S.A., Science, 254:1146-1153 (1991)).

Hepatoma-derived growth factor (HDGF), has been recently cloned (Nakamura, H. et al., J. Biol. Chem., 269(40):25143-25149 (1994)). HDGF is a heparin-binding protein which mitogenic for fibroblasts. HDGF was purified from the conditioned medium of a human hepatoma-derived cell line, HuH-7 by tritiated thymidine incorporation into Swiss 3T3 cells. HDGF has no signal peptide, yet is secreted into the medium of COS-7 cells after transfection of the cDNA clone. It is a heparin-binding protein and is ubiquitously expressed

in several tumor-derived cell lines and tissues. It is localized in the cytoplasm of hepatoma cells and has strong growth stimulating activity.

The polypeptide of the present invention has been putatively identified as an HDGF-2 polypeptide. This identification has been made as a result of amino acid sequence homology to human HDGF.

In accordance with one aspect of the present invention, there is provided a novel mature polypeptide as well as biologically active and diagnostically or therapeutically useful fragments, analogs and derivatives thereof. The polypeptide of the present invention is of human origin.

In accordance with another aspect of the present invention, there are provided isolated nucleic acid molecules, including mRNAs, DNAs, cDNAs, genomic DNAs as well as analogs and biologically active and diagnostically or therapeutically useful fragments thereof.

In accordance with yet a further aspect of the present invention, there is provided a process for producing such polypeptide by recombinant techniques comprising culturing recombinant prokaryotic and/or eukaryotic host cells, containing a nucleic acid sequence encoding the polypeptide of the present invention, under conditions promoting expression of said protein and subsequent recovery of said protein.

In accordance with yet a further aspect of the present invention, there is provided a process for utilizing such polypeptide, or polynucleotide encoding such polypeptide for screening for agonist and antagonist compounds thereto and for therapeutic purposes, for example, promoting wound healing for example as a result of burns, tissue repair and ulcers, to treat thrombosis and arteriosclerosis, to prevent neuronal damage due to neuronal disorders and promote neuronal growth, to enhance bone and periodontal regeneration, treat sunburn, to stimulate growth and/or

differentiation of bone marrow cells and corneal endothelium, and to prevent skin aging and hair loss, and to stimulate organogenesis.

In accordance with yet a further aspect of the present invention, there is also provided nucleic acid probes comprising nucleic acid molecules of sufficient length to specifically hybridize to nucleic acid sequences encoding the polypeptide of the present invention .

In accordance with yet a further aspect of the present invention, there are provided antibodies against such polypeptides.

In accordance with another aspect of the present invention, there are provided agonists which mimic the polypeptide of the present invention by binding to and activating the receptors thereto.

In accordance with yet another aspect of the present invention, there are provided antagonists to such polypeptides, which may be used to inhibit the action of such polypeptides, for example, in the treatment of restenosis after angioplasty, tumor angiogenesis, to prevent scarring and to treat hyper-vascular diseases.

In accordance with yet another aspect of the present invention, there are provided diagnostic assays for detecting diseases or susceptibility to diseases related to mutations in a nucleic acid sequence of the present invention and for detecting over-expression of the polypeptides encoded by such sequences.

In accordance with another aspect of the present invention, there is provided a process for utilizing such polypeptides, or polynucleotides encoding such polypeptides, for *in vitro* purposes related to scientific research, synthesis of DNA and manufacture of DNA vectors.

These and other aspects of the present invention should be apparent to those skilled in the art from the teachings herein.

The following drawings are illustrative of embodiments of the invention and are not meant to limit the scope of the invention as encompassed by the claims.

Figure 1 depicts the cDNA sequence and corresponding deduced amino acid sequence of HDGF-2. The standard one letter abbreviation for amino acids is used. Sequencing was performed using a 373 Automated DNA sequencer (Applied Biosystems, Inc.).

Figure 2 is an amino acid comparison between the polypeptide of the present invention (top line) and human HDGF-1 (bottom line).

In accordance with an aspect of the present invention, there is provided an isolated nucleic acid (polynucleotide) which encodes for the mature polypeptide having the deduced amino acid sequence of Figure 1 (SEQ ID NO:2) or for the mature polypeptide encoded by the cDNA of the clone deposited as ATCC Deposit No. \_\_\_\_\_ on May 24, 1995.

A polynucleotide encoding a polypeptide of the present invention may be obtained from heart, brain, and skeletal muscle. The polynucleotide of this invention was discovered in a cDNA library derived from human umbilical vein endothelial tissue. It is structurally related to the HDGF family. It contains an open reading frame encoding a protein of 249 amino acid residues. The protein exhibits the highest degree of homology to human HDGF with 23 % identity and 61 % similarity over a 201 amino acid stretch.

The polynucleotide of the present invention may be in the form of RNA or in the form of DNA, which DNA includes cDNA, genomic DNA, and synthetic DNA. The DNA may be double-stranded or single-stranded, and if single stranded may be the coding strand or non-coding (anti-sense) strand. The coding sequence which encodes the mature polypeptide may be identical to the coding sequence shown in Figure 1 (SEQ ID NO:1) or that of the deposited clone or may be a different coding sequence which coding sequence, as a result of the

redundancy or degeneracy of the genetic code, encodes the same mature polypeptide as the DNA of Figure 1 (SEQ ID NO:1) or the deposited cDNA.

The polynucleotide which encodes for the mature polypeptide of Figure 1 (SEQ ID NO:2) or for the mature polypeptide encoded by the deposited cDNA may include, but is not limited to: only the coding sequence for the mature polypeptide; the coding sequence for the mature polypeptide and additional coding sequence; the coding sequence for the mature polypeptide (and optionally additional coding sequence) and non-coding sequence, such as introns or non-coding sequence 5' and/or 3' of the coding sequence for the mature polypeptide.

Thus, the term "polynucleotide encoding a polypeptide" encompasses a polynucleotide which includes only coding sequence for the polypeptide as well as a polynucleotide which includes additional coding and/or non-coding sequence.

The present invention further relates to variants of the hereinabove described polynucleotides which encode for fragments, analogs and derivatives of the polypeptide having the deduced amino acid sequence of Figure 1 (SEQ ID NO:2) or the polypeptide encoded by the cDNA of the deposited clone. The variant of the polynucleotide may be a naturally occurring allelic variant of the polynucleotide or a non-naturally occurring variant of the polynucleotide.

Thus, the present invention includes polynucleotides encoding the same mature polypeptide as shown in Figure 1 (SEQ ID NO:2) or the same mature polypeptide encoded by the cDNA of the deposited clone as well as variants of such polynucleotides which variants encode for a fragment, derivative or analog of the polypeptide of Figure 1 or the polypeptide encoded by the cDNA of the deposited clone. Such nucleotide variants include deletion variants, substitution variants and addition or insertion variants.

As hereinabove indicated, the polynucleotide may have a coding sequence which is a naturally occurring allelic variant of the coding sequence shown in Figure 1 (SEQ ID NO:1) or of the coding sequence of the deposited clone. As known in the art, an allelic variant is an alternate form of a polynucleotide sequence which may have a substitution, deletion or addition of one or more nucleotides, which does not substantially alter the function of the encoded polypeptide.

The present invention also includes polynucleotides, wherein the coding sequence for the mature polypeptide may be fused in the same reading frame to a polynucleotide sequence which aids in expression and secretion of a polypeptide from a host cell, for example, a leader sequence which functions as a secretory sequence for controlling transport of a polypeptide from the cell. The polypeptide having a leader sequence is a preprotein and may have the leader sequence cleaved by the host cell to form the mature form of the polypeptide. The polynucleotides may also encode for a proprotein which is the mature protein plus additional 5' amino acid residues. A mature protein having a prosequence is a proprotein and is an inactive form of the protein. Once the prosequence is cleaved an active mature protein remains.

Thus, for example, the polynucleotide of the present invention may encode for a mature protein, or for a protein having a prosequence or for a protein having both a prosequence and a presequence (leader sequence).

The polynucleotides of the present invention may also have the coding sequence fused in frame to a marker sequence which allows for purification of the polypeptide of the present invention. The marker sequence may be a hexahistidine tag supplied by a pQE-9 vector to provide for purification of the mature polypeptide fused to the marker in the case of a bacterial host, or, for example, the marker sequence may be a hemagglutinin (HA) tag when a mammalian



host, e.g. COS-7 cells, is used. The HA tag corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson, I., et al., Cell, 37:767 (1984)).

The term "gene" means the segment of DNA involved in producing a polypeptide chain; it includes regions preceding and following the coding region (leader and trailer) as well as intervening sequences (introns) between individual coding segments (exons).

Fragments of the full length HDGF-2 gene may be used as a hybridization probe for a cDNA library to isolate the full length HDGF-2 gene and to isolate other genes which have a high sequence similarity to the gene or similar biological activity. Probes of this type preferably have at least 30 bases and may contain, for example, 50 or more bases. The probe may also be used to identify a cDNA clone corresponding to a full length transcript and a genomic clone or clones that contain the complete gene including regulatory and promotor regions, exons, and introns. An example of a screen comprises isolating the coding region of the gene by using the known DNA sequence to synthesize an oligonucleotide probe. Labeled oligonucleotides having a sequence complementary to that of the gene of the present invention are used to screen a library of human cDNA, genomic DNA or mRNA to determine which members of the library the probe hybridizes to.

The present invention further relates to polynucleotides which hybridize to the hereinabove-described sequences if there is at least 70%, preferably at least 90%, and more preferably at least 95% identity between the sequences. The present invention particularly relates to polynucleotides which hybridize under stringent conditions to the hereinabove-described polynucleotides. As herein used, the term "stringent conditions" means hybridization will occur only if there is at least 95% and preferably at least 97% identity between the sequences. The polynucleotides

which hybridize to the hereinabove described polynucleotides in a preferred embodiment encode polypeptides which either retain substantially the same biological function or activity as the mature polypeptide encoded by the cDNAs of Figure 1 (SEQ ID NO:1) or the deposited cDNA(s).

Alternatively, the polynucleotide may have at least 20 bases, preferably 30 bases, and more preferably at least 50 bases which hybridize to a polynucleotide of the present invention and which has an identity thereto, as hereinabove described, and which may or may not retain activity. For example, such polynucleotides may be employed as probes for the polynucleotide of SEQ ID NO:1, for example, for recovery of the polynucleotide or as a diagnostic probe or as a PCR primer.

Thus, the present invention is directed to polynucleotides having at least a 70% identity, preferably at least 90% and more preferably at least a 95% identity to a polynucleotide which encodes the polypeptide of SEQ ID NO:2 as well as fragments thereof, which fragments have at least 30 bases and preferably at least 50 bases and to polypeptides encoded by such polynucleotides.

The deposit(s) referred to herein will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Micro-organisms for purposes of Patent Procedure. These deposits are provided merely as convenience to those of skill in the art and are not an admission that a deposit is required under 35 U.S.C. §112. The sequence of the polynucleotides contained in the deposited materials, as well as the amino acid sequence of the polypeptides encoded thereby, are incorporated herein by reference and are controlling in the event of any conflict with any description of sequences herein. A license may be required to make, use or sell the deposited materials, and no such license is hereby granted.

The present invention further relates to an HDGF-2 polypeptide which has the deduced amino acid sequence of Figure 1 (SEQ ID NO:2) or which has the amino acid sequence encoded by the deposited cDNA, as well as fragments, analogs and derivatives of such polypeptide.

The terms "fragment," "derivative" and "analog" when referring to the polypeptide of Figure 1 (SEQ ID NO:2) or that encoded by the deposited cDNA, means a polypeptide which retains essentially the same biological function or activity as such polypeptide. Thus, an analog includes a proprotein which can be activated by cleavage of the proprotein portion to produce an active mature polypeptide.

The polypeptide of the present invention may be a recombinant polypeptide, a natural polypeptide or a synthetic polypeptide, preferably a recombinant polypeptide.

The fragment, derivative or analog of the polypeptide of Figure 1 (SEQ ID NO:2) or that encoded by the deposited cDNA may be (i) one in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one encoded by the genetic code, or (ii) one in which one or more of the amino acid residues includes a substituent group, or (iii) one in which the mature polypeptide is fused with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol), or (iv) one in which the additional amino acids are fused to the mature polypeptide and employed for purification of the mature polypeptide. Such fragments, derivatives and analogs are deemed to be within the scope of those skilled in the art from the teachings herein.

The polypeptides and polynucleotides of the present invention are preferably provided in an isolated form, and preferably are purified to homogeneity.

The term "isolated" means that the material is removed from its original environment (e.g., the natural environment if it is naturally occurring). For example, a naturally-occurring polynucleotide or polypeptide present in a living animal is not isolated, but the same polynucleotide or polypeptide, separated from some or all of the coexisting materials in the natural system, is isolated. Such polynucleotides could be part of a vector and/or such polynucleotides or polypeptides could be part of a composition, and still be isolated in that such vector or composition is not part of its natural environment.

The polypeptides of the present invention include the polypeptide of SEQ ID NO:2 (in particular the mature polypeptide) as well as polypeptides which have at least 70% similarity (preferably at least a 70% identity) to the polypeptide of SEQ ID NO:2 and more preferably at least a 90% similarity (more preferably at least a 90% identity) to the polypeptide of SEQ ID NO:2 and still more preferably at least a 95% similarity (still more preferably at least a 95% identity) to the polypeptide of SEQ ID NO:2 and also include portions of such polypeptides with such portion of the polypeptide generally containing at least 30 amino acids and more preferably at least 50 amino acids.

As known in the art "similarity" between two polypeptides is determined by comparing the amino acid sequence and its conserved amino acid substitutes of one polypeptide to the sequence of a second polypeptide.

Fragments or portions of the polypeptides of the present invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, the fragments may be employed as intermediates for producing the full-length polypeptides. Fragments or portions of the polynucleotides of the present invention may be used to synthesize full-length polynucleotides of the present invention.

The present invention also relates to vectors which include polynucleotides of the present invention, host cells which are genetically engineered with vectors of the invention and the production of polypeptides of the invention by recombinant techniques.

Host cells are genetically engineered (transduced or transformed or transfected) with the vectors of this invention which may be, for example, a cloning vector or an expression vector. The vector may be, for example, in the form of a plasmid, a viral particle, a phage, etc. The engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants or amplifying the genes of the present invention. The culture conditions, such as temperature, pH and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

The polynucleotides of the present invention may be employed for producing polypeptides by recombinant techniques. Thus, for example, the polynucleotide may be included in any one of a variety of expression vectors for expressing a polypeptide. Such vectors include chromosomal, nonchromosomal and synthetic DNA sequences, e.g., derivatives of SV40; bacterial plasmids; phage DNA; baculovirus; yeast plasmids; vectors derived from combinations of plasmids and phage DNA, viral DNA such as vaccinia, adenovirus, fowl pox virus, and pseudorabies. However, any other vector may be used as long as it is replicable and viable in the host.

The appropriate DNA sequence may be inserted into the vector by a variety of procedures. In general, the DNA sequence is inserted into an appropriate restriction endonuclease site(s) by procedures known in the art. Such procedures and others are deemed to be within the scope of those skilled in the art.

The DNA sequence in the expression vector is operatively linked to an appropriate expression control sequence(s) (promoter) to direct mRNA synthesis. As representative examples of such promoters, there may be mentioned: LTR or SV40 promoter, the E. coli. lac or trp, the phage lambda P<sub>L</sub> promoter and other promoters known to control expression of genes in prokaryotic or eukaryotic cells or their viruses. The expression vector also contains a ribosome binding site for translation initiation and a transcription terminator. The vector may also include appropriate sequences for amplifying expression.

In addition, the expression vectors preferably contain one or more selectable marker genes to provide a phenotypic trait for selection of transformed host cells such as dihydrofolate reductase or neomycin resistance for eukaryotic cell culture, or such as tetracycline or ampicillin resistance in E. coli.

The vector containing the appropriate DNA sequence as hereinabove described, as well as an appropriate promoter or control sequence, may be employed to transform an appropriate host to permit the host to express the protein.

As representative examples of appropriate hosts, there may be mentioned: bacterial cells, such as E. coli, Streptomyces, Salmonella typhimurium; fungal cells, such as yeast; insect cells such as Drosophila S2 and Spodoptera Sf9; animal cells such as CHO, COS or Bowes melanoma; adenoviruses; plant cells, etc. The selection of an appropriate host is deemed to be within the scope of those skilled in the art from the teachings herein.

More particularly, the present invention also includes recombinant constructs comprising one or more of the sequences as broadly described above. The constructs comprise a vector, such as a plasmid or viral vector, into which a sequence of the invention has been inserted, in a forward or reverse orientation. In a preferred aspect of this

embodiment, the construct further comprises regulatory sequences, including, for example, a promoter, operably linked to the sequence. Large numbers of suitable vectors and promoters are known to those of skill in the art, and are commercially available. The following vectors are provided by way of example; Bacterial: pQE70, pQE60, pQE-9 (Qiagen), pBS, pD10, phagescript, psiX174, pbluescript SK, pbsks, pNH8A, pNH16a, pNH18A, pNH46A (Stratagene); ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 (Pharmacia); Eukaryotic: pWLNEO, pSV2CAT, pOG44, pXT1, pSG (Stratagene) pSVK3, pBPV, pMSG, pSVL (Pharmacia). However, any other plasmid or vector may be used as long as they are replicable and viable in the host.

Promoter regions can be selected from any desired gene using CAT (chloramphenicol transferase) vectors or other vectors with selectable markers. Two appropriate vectors are pKK232-8 and pCM7. Particular named bacterial promoters include lacI, lacZ, T3, T7, gpt, lambda P<sub>R</sub>, P<sub>L</sub> and trp. Eukaryotic promoters include CMV immediate early, HSV thymidine kinase, early and late SV40, LTRs from retrovirus, and mouse metallothionein-I. Selection of the appropriate vector and promoter is well within the level of ordinary skill in the art.

In a further embodiment, the present invention relates to host cells containing the above-described constructs. The host cell can be a higher eukaryotic cell, such as a mammalian cell, or a lower eukaryotic cell, such as a yeast cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-Dextran mediated transfection, or electroporation (Davis, L., Dibner, M., Battey, I., Basic Methods in Molecular Biology, (1986)).

The constructs in host cells can be used in a conventional manner to produce the gene product encoded by

the recombinant sequence. Alternatively, the polypeptides of the invention can be synthetically produced by conventional peptide synthesizers.

Mature proteins can be expressed in mammalian cells, yeast, bacteria, or other cells under the control of appropriate promoters. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention. Appropriate cloning and expression vectors for use with prokaryotic and eukaryotic hosts are described by Sambrook, et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor, N.Y., (1989), the disclosure of which is hereby incorporated by reference.

Transcription of the DNA encoding the polypeptides of the present invention by higher eukaryotes is increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp that act on a promoter to increase its transcription. Examples include the SV40 enhancer on the late side of the replication origin bp 100 to 270, a cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers.

Generally, recombinant expression vectors will include origins of replication and selectable markers permitting transformation of the host cell, e.g., the ampicillin resistance gene of *E. coli* and *S. cerevisiae* TRP1 gene, and a promoter derived from a highly-expressed gene to direct transcription of a downstream structural sequence. Such promoters can be derived from operons encoding glycolytic enzymes such as 3-phosphoglycerate kinase (PGK),  $\alpha$ -factor, acid phosphatase, or heat shock proteins, among others. The heterologous structural sequence is assembled in appropriate phase with translation initiation and termination sequences. Optionally, the heterologous sequence can encode a fusion protein including an N-terminal identification peptide



imparting desired characteristics, e.g., stabilization or simplified purification of expressed recombinant product.

Useful expression vectors for bacterial use are constructed by inserting a structural DNA sequence encoding a desired protein together with suitable translation initiation and termination signals in operable reading phase with a functional promoter. The vector will comprise one or more phenotypic selectable markers and an origin of replication to ensure maintenance of the vector and to, if desirable, provide amplification within the host. Suitable prokaryotic hosts for transformation include E. coli, Bacillus subtilis, Salmonella typhimurium and various species within the genera Pseudomonas, Streptomyces, and Staphylococcus, although others may also be employed as a matter of choice.

As a representative but nonlimiting example, useful expression vectors for bacterial use can comprise a selectable marker and bacterial origin of replication derived from commercially available plasmids comprising genetic elements of the well known cloning vector pBR322 (ATCC 37017). Such commercial vectors include, for example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and GEM1 (Promega Biotec, Madison, WI, USA). These pBR322 "backbone" sections are combined with an appropriate promoter and the structural sequence to be expressed.

Following transformation of a suitable host strain and growth of the host strain to an appropriate cell density, the selected promoter is induced by appropriate means (e.g., temperature shift or chemical induction) and cells are cultured for an additional period.

Cells are typically harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification.

Microbial cells employed in expression of proteins can be disrupted by any convenient method, including freeze-thaw

cycling, sonication, mechanical disruption, or use of cell lysing agents, such methods are well known to those skilled in the art.

Various mammalian cell culture systems can also be employed to express recombinant protein. Examples of mammalian expression systems include the COS-7 lines of monkey kidney fibroblasts, described by Gluzman, Cell, 23:175 (1981), and other cell lines capable of expressing a compatible vector, for example, the C127, 3T3, CHO, HeLa and BHK cell lines. Mammalian expression vectors will comprise an origin of replication, a suitable promoter and enhancer, and also any necessary ribosome binding sites, polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking nontranscribed sequences. DNA sequences derived from the SV40 splice, and polyadenylation sites may be used to provide the required nontranscribed genetic elements.

The HDGF-2 polypeptide can be recovered and purified from recombinant cell cultures by methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Protein refolding steps can be used, as necessary, in completing configuration of the mature protein. Finally, high performance liquid chromatography (HPLC) can be employed for final purification steps.

The polypeptides of the present invention may be a naturally purified product, or a product of chemical synthetic procedures, or produced by recombinant techniques from a prokaryotic or eukaryotic host (for example, by bacterial, yeast, higher plant, insect and mammalian cells in culture). Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated.

Polypeptides of the invention may also include an initial methionine amino acid residue.

The polypeptides of the present invention, as a result of angiogenic ability, stimulate vascular endothelial cell growth, and may be employed in treatment for stimulating re-vascularization of ischaemic tissues due to various disease conditions such as thrombosis, arteriosclerosis, and other cardiovascular conditions.

These polypeptides may also be employed to stimulate mesodermal induction and limb regeneration in early embryos.

The polypeptides may also be employed for promoting healing in wounds due to injuries, burns, surgery, and ulcers since they are mitogenic to various cells of different origins, such as fibroblast cells and skeletal muscle cells, and therefore, facilitate the repair or replacement of damaged or diseased tissue.

The polypeptides of the present invention may also be employed stimulate neuronal growth and to treat and prevent neuronal damage which occurs in certain neuronal disorders or neuro-degenerative conditions such as Alzheimer's disease, Parkinson's disease, and AIDS-related complex.

The polypeptides may be employed to stimulate chondrocyte growth, therefore, they may be employed to enhance bone and periodontal regeneration and aid in tissue transplants or bone grafts.

The polypeptides of the present invention may be also be employed to prevent skin aging due to sunburn by stimulating keratinocyte growth.

The polypeptides may also be employed for preventing hair loss by activating hair-forming cells and promoting melanocyte growth.

The polypeptides of the present invention may be employed to stimulate growth and differentiation of hematopoietic cells and bone marrow cells.

The polypeptides may also be employed to maintain organs before transplantation or for supporting cell culture of primary tissues.

The polypeptide of the present invention may also be employed for inducing tissue of mesodermal origin to differentiate in early embryos.

The polynucleotides and polypeptides of the present invention may be employed as research reagents and materials for discovery of treatments and diagnostics to human disease.

This invention provides a method for identification of the receptor for the polypeptide of the present invention. The gene encoding the receptor can be identified by numerous methods known to those of skill in the art, for example, ligand panning and FACS sorting (Coligan, et al., Current Protocols in Immun., 1(2), Chapter 5, (1991)). Preferably, expression cloning is employed wherein polyadenylated RNA is prepared from a cell responsive to the HDGF-2 polypeptide, and a cDNA library created from this RNA is divided into pools and used to transfect COS cells or other cells that are not responsive. Transfected cells which are grown on glass slides are exposed to labeled HDGF-2. The HDGF-2 polypeptide can be labeled by a variety of means including iodination or inclusion of a recognition site for a site-specific protein kinase. Following fixation and incubation, the slides are subjected to auto-radiographic analysis. Positive pools are identified and sub-pools are prepared and re-transfected using an iterative sub-pooling and re-screening process, eventually yielding a single clone that encodes the putative receptor. As an alternative approach for receptor identification, labeled ligand can be photoaffinity linked with cell membrane or extract preparations that express the receptor molecule. Cross-linked material is resolved by PAGE and exposed to X-ray film. The labeled complex containing the ligand-receptor can be excised, resolved into peptide fragments, and subjected to protein microsequencing. The

amino acid sequence obtained from microsequencing would be used to design a set of degenerate oligonucleotide probes to screen a cDNA library to identify the gene encoding the putative receptor.

This invention provides a method of screening compounds to identify those which bind to and activate the HDGF-2 receptor (agonists) or bind to and inhibit the HDGF-1 receptor (antagonists). As an example, a competition assay may be employed wherein a mammalian cell or membrane preparation expressing the HDGF-2 receptor is incubated with labeled HDGF-2 and the compound to be tested. The ability of the compound to compete for the HDGF-2 receptors can be determined by using liquid scintillation counting to determine the amount of bound HDGF-2 polypeptide in the presence and absence of the compound.

Alternatively, agonist compounds may be identified by detecting the response of a known second messenger system following interaction of a compound to be tested and the HDGF-2 receptor is measured by labeling the compound with radioactivity and contacting it with a cell expressing the HDGF-2 receptor. Such second messenger systems include but are not limited to, cAMP guanylate cyclase, ion channels or phosphoinositide hydrolysis. Antagonist compounds may be determined using this assay system wherein antagonists are determined to bind to the receptor but not elicit a second messenger response to, therefore, effectively block HDGF-2 polypeptide from its receptor.

Methods to prepare cells for the above-described assays, comprises transfecting a cell population (one presumed not to contain the receptor) with the appropriate vector containing DNA encoding the HDGF-2 receptor, such that the cell will now express the receptor. A suitable response system is obtained by transfection of the DNA into a suitable host containing the desired second messenger pathways including cAMP, ion channels, phosphoinositide kinase, or calcium response. Such

a transfection system provides a response system to analyze the activity of various compounds and polypeptides exposed to the cell.

In a specific assay, growth stimulating activity of HDGF-2 can be assaying in the presence of potential compounds by measuring [<sup>3</sup>H]thymidine incorporation into Weiss albino mouse 3T3 cells as per Nakamura, H., et al., Clin. Chim. Acta, 183:273-284 (1989) which is hereby incorporated by reference in its entirety.

Examples of antagonist compounds include antibodies which are immunoreactive with various critical positions on the HDGF-2 receptor, and bind to the receptor and block it from HDGF-2 polypeptide. Antibodies include anti-idiotypic antibodies which recognize unique determinants generally associated with the antigen-binding site of an antibody. Antibodies of this type may also be prepared against the HDGF-2 polypeptide itself to bind thereto and prevent it from interacting with its receptor.

Oligopeptides which bind to the HDGF-2 receptor in competition with HDGF-2 itself but which do not elicit a second messenger response, may also be used as antagonist compounds. Examples of oligopeptides include small molecules, for example, small peptides or peptide-like molecules. Oligopeptides may also bind to the active site of HDGF-2 to block interaction of HDGF-2 with its receptor.

Antisense construct prepared using antisense technology may also be employed as antagonist compounds to prevent product of the HDGF-2 polypeptide. Antisense technology can be used to control gene expression through triple-helix formation or antisense DNA or RNA, both of which methods are based on binding of a polynucleotide to DNA or RNA. For example, the 5' coding portion of the polynucleotide sequence, which encodes for the mature polypeptides of the present invention, is used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length.

A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription (triple helix - see Lee et al., Nucl. Acids Res., 6:3073 (1979); Cooney et al, Science, 241:456 (1988); and Dervan et al., Science, 251: 1360 (1991)), thereby preventing transcription and the production of HDGF-2. The antisense RNA oligonucleotide hybridizes to the mRNA *in vivo* and blocks translation of the mRNA molecule into HDGF-2 polypeptide (Antisense - Okano, J. Neurochem., 56:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988)). The oligonucleotides described above can also be delivered to cells such that the antisense RNA or DNA may be expressed *in vivo* to inhibit production of HDGF-2.

The antagonists may be employed to inhibit the cell growth and proliferation effects of the polypeptides of the present invention on neoplastic cells and tissues, i.e. stimulation of angiogenesis of tumors, and, therefore, retard or prevent abnormal cellular growth and proliferation, for example, in tumor formation or growth.

The antagonists may also be employed to prevent hyper-vascular diseases, and prevent the proliferation of epithelial lens cells after extracapsular cataract surgery.

Prevention of the mitogenic activity of the polypeptides of the present invention may also be desirable in cases such as restenosis after balloon angioplasty.

The antagonists may also be employed to prevent inflammation and the growth of scar tissue during wound healing.

The antagonists may be employed in a composition with a pharmaceutically acceptable carrier, e.g., as hereinafter described.

The polypeptides of the present invention and agonist and antagonist compounds may be employed in combination with a suitable pharmaceutical carrier. Such compositions comprise a therapeutically effective amount of the

polypeptide or compound, and a pharmaceutically acceptable carrier or excipient. Such a carrier includes but is not limited to saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The formulation should suit the mode of administration.

The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration. In addition, the polypeptides and compounds of the present invention may be employed in conjunction with other therapeutic compounds.

The pharmaceutical compositions may be administered in a convenient manner such as by the topical, intravenous, intraperitoneal, intramuscular, subcutaneous or intradermal routes. The pharmaceutical compositions are administered in an amount which is effective for treating and/or prophylaxis of the specific indication. In general, they are administered in an amount of at least about 10  $\mu\text{g/kg}$  body weight and in most cases they will be administered in an amount not in excess of about 8 mg/Kg body weight per day. In most cases, the dosage is from about 10  $\mu\text{g/kg}$  to about 1 mg/kg body weight daily, taking into account the routes of administration, symptoms, etc.

The HDGF-2 polypeptides and agonists and antagonists which are polypeptides may also be employed in accordance with the present invention by expression of such polypeptides *in vivo*, which is often referred to as "gene therapy."

Thus, for example, cells from a patient may be engineered with a polynucleotide (DNA or RNA) encoding a polypeptide *ex vivo*, with the engineered cells then being



provided to a patient to be treated with the polypeptide. Such methods are well-known in the art and are apparent from the teachings herein. For example, cells may be engineered by the use of a retroviral plasmid vector containing RNA encoding a polypeptide of the present invention.

Similarly, cells may be engineered *in vivo* for expression of a polypeptide *in vivo* by, for example, procedures known in the art. For example, a packaging cell is transduced with a retroviral plasmid vector containing RNA encoding a polypeptide of the present invention such that the packaging cell now produces infectious viral particles containing the gene of interest. These producer cells may be administered to a patient for engineering cells *in vivo* and expression of the polypeptide *in vivo*. These and other methods for administering a polypeptide of the present invention by such method should be apparent to those skilled in the art from the teachings of the present invention.

Retroviruses from which the retroviral plasmid vectors hereinabove mentioned may be derived include, but are not limited to, Moloney Murine Leukemia Virus, spleen necrosis virus, retroviruses such as Rous Sarcoma Virus, Harvey Sarcoma Virus, avian leukosis virus, gibbon ape leukemia virus, human immunodeficiency virus, adenovirus, Myeloproliferative Sarcoma Virus, and mammary tumor virus. In one embodiment, the retroviral plasmid vector is derived from Moloney Murine Leukemia Virus.

The vector includes one or more promoters. Suitable promoters which may be employed include, but are not limited to, the retroviral LTR; the SV40 promoter; and the human cytomegalovirus (CMV) promoter described in Miller, et al., Biotechniques, Vol. 7, No. 9, 980-990 (1989), or any other promoter (e.g., cellular promoters such as eukaryotic cellular promoters including, but not limited to, the histone, pol III, and  $\beta$ -actin promoters). Other viral promoters which may be employed include, but are not limited

to, adenovirus promoters, thymidine kinase (TK) promoters, and B19 parvovirus promoters. The selection of a suitable promoter will be apparent to those skilled in the art from the teachings contained herein.

The nucleic acid sequence encoding the polypeptide of the present invention is under the control of a suitable promoter. Suitable promoters which may be employed include, but are not limited to, adenoviral promoters, such as the adenoviral major late promoter; or heterologous promoters, such as the cytomegalovirus (CMV) promoter; the respiratory syncytial virus (RSV) promoter; inducible promoters, such as the MMT promoter, the metallothionein promoter; heat shock promoters; the albumin promoter; the ApoA1 promoter; human globin promoters; viral thymidine kinase promoters, such as the Herpes Simplex thymidine kinase promoter; retroviral LTRs (including the modified retroviral LTRs hereinabove described); the  $\beta$ -actin promoter; and human growth hormone promoters. The promoter also may be the native promoter which controls the gene encoding the polypeptide.

The retroviral plasmid vector is employed to transduce packaging cell lines to form producer cell lines. Examples of packaging cells which may be transfected include, but are not limited to, the PE501, PA317,  $\psi$ -2,  $\psi$ -AM, PA12, T19-14X, VT-19-17-H2,  $\psi$ CRE,  $\psi$ CRIP, GP+E-86, GP+envAm12, and DAN cell lines as described in Miller, Human Gene Therapy, Vol. 1, pgs. 5-14 (1990), which is incorporated herein by reference in its entirety. The vector may transduce the packaging cells through any means known in the art. Such means include, but are not limited to, electroporation, the use of liposomes, and CaPO<sub>4</sub> precipitation. In one alternative, the retroviral plasmid vector may be encapsulated into a liposome, or coupled to a lipid, and then administered to a host.

The producer cell line generates infectious retroviral vector particles which include the nucleic acid sequence(s)

encoding the polypeptides. Such retroviral vector particles then may be employed, to transduce eukaryotic cells, either *in vitro* or *in vivo*. The transduced eukaryotic cells will express the nucleic acid sequence(s) encoding the polypeptide. Eukaryotic cells which may be transduced include, but are not limited to, embryonic stem cells, embryonic carcinoma cells, as well as hematopoietic stem cells, hepatocytes, fibroblasts, myoblasts, keratinocytes, endothelial cells, and bronchial epithelial cells.

This invention is also related to the use of the HDGF-2 gene as a diagnostic. Detection of a mutation in the nucleic acid sequences encoding HDGF-2 will allow a diagnosis of a disease or a susceptibility to a disease which results from underexpression of HDGF-2.

Individuals carrying mutations in the human HDGF-2 gene may be detected at the DNA level by a variety of techniques. Nucleic acids for diagnosis may be obtained from a patient's cells, such as from blood, urine, saliva, tissue biopsy and autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR (Saiki et al., *Nature*, 324:163-166 (1986)) prior to analysis. RNA or cDNA may also be used for the same purpose. As an example, PCR primers complementary to the nucleic acid encoding the polypeptide of the present invention can be used to identify and analyze HDGF-2 mutations. For example, deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to radiolabeled HDGF-2 RNA or alternatively, radiolabeled HDGF-2 antisense DNA sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase A digestion or by differences in melting temperatures.

Sequence differences between the reference gene and genes having mutations may be revealed by the direct DNA

sequencing method. In addition, cloned DNA segments may be employed as probes to detect specific DNA segments. The sensitivity of this method is greatly enhanced when combined with PCR. For example, a sequencing primer is used with double-stranded PCR product or a single-stranded template molecule generated by a modified PCR. The sequence determination is performed by conventional procedures with radiolabeled nucleotide or by automatic sequencing procedures with fluorescent-tags.

Genetic testing based on DNA sequence differences may be achieved by detection of alteration in electrophoretic mobility of DNA fragments in gels with or without denaturing agents. Small sequence deletions and insertions can be visualized by high resolution gel electrophoresis. DNA fragments of different sequences may be distinguished on denaturing formamide gradient gels in which the mobilities of different DNA fragments are retarded in the gel at different positions according to their specific melting or partial melting temperatures (see, e.g., Myers et al., Science, 230:1242 (1985)).

Sequence changes at specific locations may also be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method (e.g., Cotton et al., PNAS, USA, 85:4397-4401 (1985)).

Thus, the detection of a specific DNA sequence may be achieved by methods such as hybridization, RNase protection, chemical cleavage, direct DNA sequencing or the use of restriction enzymes, (e.g., Restriction Fragment Length Polymorphisms (RFLP)) and Southern blotting of genomic DNA.

In addition to more conventional gel-electrophoresis and DNA sequencing, mutations can also be detected by *in situ* analysis.

The present invention also relates to a diagnostic assay for detecting altered levels of HDGF-2 protein in various tissues since an over-expression of the proteins compared to

normal control tissue samples can detect the presence of abnormal cellular differentiation and growth, for example, as occurs in neoplasia. Assays used to detect levels of HDGF-2 protein in a sample derived from a host are well-known to those of skill in the art and include radioimmunoassays, competitive-binding assays, Western Blot analysis and preferably an ELISA assay. An ELISA assay initially comprises preparing an antibody specific to the HDGF-2 antigen, preferably a monoclonal antibody. In addition a reporter antibody is prepared against the monoclonal antibody. To the reporter antibody is attached a detectable reagent such as radioactivity, fluorescence or in this example a horseradish peroxidase enzyme. A sample is now removed from a host and incubated on a solid support, e.g. a polystyrene dish, that binds the proteins in the sample. Any free protein binding sites on the dish are then covered by incubating with a non-specific protein such as bovine serum albumin. Next, the monoclonal antibody is incubated in the dish during which time the monoclonal antibodies attach to any HDGF-2 proteins attached to the polystyrene dish. All unbound monoclonal antibody is washed out with buffer. The reporter antibody linked to horseradish peroxidase is now placed in the dish resulting in binding of the reporter antibody to any monoclonal antibody bound to HDGF-2. Unattached reporter antibody is then washed out. Peroxidase substrates are then added to the dish and the amount of color developed in a given time period is a measurement of the amount of HDGF-2 protein present in a given volume of patient sample when compared against a standard curve.

A competition assay may be employed wherein antibodies specific to HDGF-2 are attached to a solid support and labeled HDGF-2 and a sample derived from the host are passed over the solid support and the amount of label detected attached to the solid support can be correlated to a quantity of HDGF-2 in the sample.

The sequences of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome. Moreover, there is a current need for identifying particular sites on the chromosome. Few chromosome marking reagents based on actual sequence data (repeat polymorphisms) are presently available for marking chromosomal location. The mapping of DNAs to chromosomes according to the present invention is an important first step in correlating those sequences with genes associated with disease.

Briefly, sequences can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp) from the cDNA. Computer analysis of the 3' untranslated region of the gene is used to rapidly select primers that do not span more than one exon in the genomic DNA, thus complicating the amplification process. These primers are then used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the primer will yield an amplified fragment.

PCR mapping of somatic cell hybrids is a rapid procedure for assigning a particular DNA to a particular chromosome. Using the present invention with the same oligonucleotide primers, sublocalization can be achieved with panels of fragments from specific chromosomes or pools of large genomic clones in an analogous manner. Other mapping strategies that can similarly be used to map to its chromosome include *in situ* hybridization, prescreening with labeled flow-sorted chromosomes and preselection by hybridization to construct chromosome specific-cDNA libraries.

Fluorescence *in situ* hybridization (FISH) of a cDNA clone to a metaphase chromosomal spread can be used to provide a precise chromosomal location in one step. This technique can be used with cDNA having at least 50 or 60 bases. For a review of this technique, see Verma et al.,

Human Chromosomes: a Manual of Basic Techniques, Pergamon Press, New York (1988).

Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes).

Next, it is necessary to determine the differences in the cDNA or genomic sequence between affected and unaffected individuals. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

With current resolution of physical mapping and genetic mapping techniques, a cDNA precisely localized to a chromosomal region associated with the disease could be one of between 50 and 500 potential causative genes. (This assumes 1 megabase mapping resolution and one gene per 20 kb).

The polypeptides, their fragments or other derivatives, or analogs thereof, or cells expressing them can be used as an immunogen to produce antibodies thereto. These antibodies can be, for example, polyclonal or monoclonal antibodies. The present invention also includes chimeric, single chain, and humanized antibodies, as well as Fab fragments, or the product of an Fab expression library. Various procedures known in the art may be used for the production of such antibodies and fragments.

Antibodies generated against the polypeptides corresponding to a sequence of the present invention can be obtained by direct injection of the polypeptides into an

animal or by administering the polypeptides to an animal, preferably a nonhuman. The antibody so obtained will then bind the polypeptides itself. In this manner, even a sequence encoding only a fragment of the polypeptides can be used to generate antibodies binding the whole native polypeptides. Such antibodies can then be used to isolate the polypeptide from tissue expressing that polypeptide.

For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler and Milstein, 1975, *Nature*, 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor et al., 1983, *Immunology Today* 4:72), and the EBV-hybridoma technique to produce human monoclonal antibodies (Cole, et al., 1985, in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96).

Techniques described for the production of single chain antibodies (U.S. Patent 4,946,778) can be adapted to produce single chain antibodies to immunogenic polypeptide products of this invention. Also, transgenic mice may be used to express humanized antibodies to immunogenic polypeptide products of this invention.

The present invention will be further described with reference to the following examples; however, it is to be understood that the present invention is not limited to such examples. All parts or amounts, unless otherwise specified, are by weight.

In order to facilitate understanding of the following examples certain frequently occurring methods and/or terms will be described.

"Plasmids" are designated by a lower case p preceded and/or followed by capital letters and/or numbers. The starting plasmids herein are either commercially available, publicly available on an unrestricted basis, or can be constructed from available plasmids in accord with published



procedures. In addition, equivalent plasmids to those described are known in the art and will be apparent to the ordinarily skilled artisan.

"Digestion" of DNA refers to catalytic cleavage of the DNA with a restriction enzyme that acts only at certain sequences in the DNA. The various restriction enzymes used herein are commercially available and their reaction conditions, cofactors and other requirements were used as would be known to the ordinarily skilled artisan. For analytical purposes, typically 1  $\mu$ g of plasmid or DNA fragment is used with about 2 units of enzyme in about 20  $\mu$ l of buffer solution. For the purpose of isolating DNA fragments for plasmid construction, typically 5 to 50  $\mu$ g of DNA are digested with 20 to 250 units of enzyme in a larger volume. Appropriate buffers and substrate amounts for particular restriction enzymes are specified by the manufacturer. Incubation times of about 1 hour at 37°C are ordinarily used, but may vary in accordance with the supplier's instructions. After digestion the reaction is electrophoresed directly on a polyacrylamide gel to isolate the desired fragment.

Size separation of the cleaved fragments is performed using 8 percent polyacrylamide gel described by Goeddel, D. et al., *Nucleic Acids Res.*, 8:4057 (1980).

"Oligonucleotides" refers to either a single stranded polydeoxynucleotide or two complementary polydeoxynucleotide strands which may be chemically synthesized. Such synthetic oligonucleotides have no 5' phosphate and thus will not ligate to another oligonucleotide without adding a phosphate with an ATP in the presence of a kinase. A synthetic oligonucleotide will ligate to a fragment that has not been dephosphorylated.

"Ligation" refers to the process of forming phosphodiester bonds between two double stranded nucleic acid fragments (Maniatis, T., et al., *Id.*, p. 146). Unless

otherwise provided, ligation may be accomplished using known buffers and conditions with 10 units of T4 DNA ligase ("ligase") per 0.5  $\mu$ g of approximately equimolar amounts of the DNA fragments to be ligated.

Unless otherwise stated, transformation was performed as described in the method of Graham, F. and Van der Eb, A., Virology, 52:456-457 (1973).

#### Example 1

##### Bacterial Expression and Purification of HDGF-2 proteins

The DNA sequence encoding HDGF-2, ATCC # \_\_, is initially amplified using PCR oligonucleotide primers corresponding to the 5' sequences of the protein and the vector sequences 3' to the gene. Additional nucleotides corresponding to the gene are added to the 5' and 3' sequences respectively. The HDGF-2 5' oligonucleotide primer has the sequence 5' ACGTGGATCCGCGCTGTGAGTCTGCGGCTCGGC 3' (SEQ ID NO:3) contains a BamHI restriction enzyme site. The 3' sequence 5' CAACAAGCTTTTCACCTAGGAAGAAGGAGGTCTTCA 3' (SEQ ID NO:4) contains complementary sequences to a HindIII site and is followed by TGF $\alpha$ -HII coding sequence.

The restriction enzyme sites correspond to the restriction enzyme sites on the bacterial expression vector pQE-9 (Qiagen, Inc. Chatsworth, CA 91311). pQE-9 encodes antibiotic resistance (Amp<sup>r</sup>), a bacterial origin of replication (ori), an IPTG-regulatable promoter operator (P/O), a ribosome binding site (RBS), a 6-His tag and restriction enzyme sites. pQE-9 was then digested with BamHI and HindIII. The amplified sequences are ligated into pQE-9, and are inserted in frame with the sequence encoding for the histidine tag and the RBS. The ligation mixture is then used to transform E. coli strain M15/rep 4 (Qiagen, Inc.) by the procedure described in Sambrook, J. et al., Molecular Cloning: A Laboratory Manual, Cold Spring Laboratory Press, (1989). M15/rep4 contains multiple copies of the plasmid

pREP4, which expresses the lacI repressor and also confers kanamycin resistance (Kan<sup>r</sup>). Transformants are identified by their ability to grow on LB plates and ampicillin/kanamycin resistant colonies were selected. Plasmid DNA is isolated and confirmed by restriction analysis. Clones containing the desired constructs are grown overnight (O/N) in liquid culture in LB media supplemented with both Amp (100 ug/ml) and Kan (25 ug/ml). The O/N culture is used to inoculate a large culture at a ratio of 1:100 to 1:250. The cells are grown to an optical density 600 (O.D.<sup>600</sup>) of between 0.4 and 0.6. IPTG ("Isopropyl-B-D-thiogalacto pyranoside") is then added to a final concentration of 1 mM. IPTG induces by inactivating the lacI repressor, clearing the P/O leading to increased gene expression. Cells are grown an extra 3 to 4 hours. Cells are then harvested by centrifugation. The cell pellet is solubilized in the chaotropic agent 6 Molar Guanidine HCl. After clarification, solubilized HDGF-2 is purified from this solution by chromatography on a Nickel-Chelate column under conditions that allow for tight binding by proteins containing the 6-His tag (Hochuli, E. et al., J. Chromatography 411:177-184 (1984)). The proteins are eluted from the column in 6 molar guanidine HCl pH 5.0 and for the purpose of renaturation adjusted to 3 molar guanidine HCl, 100mM sodium phosphate, 10 mmolar glutathione (reduced) and 2 mmolar glutathione (oxidized). After incubation in this solution for 12 hours the proteins are dialyzed to 10 mmolar sodium phosphate.

#### Example 2

##### Expression and Purification of Chemokine HDGF-2 using a baculovirus expression system.

The DNA sequence encoding the full length HDGF-2 protein, ATCC # \_\_\_\_\_, is amplified using PCR oligonucleotide primers corresponding to the 5' and 3' sequences of the gene:

The 5' primer has the sequence 5' ACGAGGATCCGCCAT

CATGGCGGCTGTGAGTCTGCGGCTCG 3' (SEQ ID NO:5) and contains a BamHI restriction enzyme site (in bold) followed by 6 nucleotides resembling an efficient signal for the initiation of translation in eukaryotic cells (Kozak, M., J. Mol. Biol., 196:947-950 (1987) and followed by 25 nucleotides of the HDGF-2 gene (the initiation codon for translation "ATG" is underlined).

The 3' primer has the sequence 5' GCATGGTACCTCACCT AGGAAGAAGGAGGTCTTCAC 3' (SEQ ID NO:6) and contains the cleavage site for the restriction endonuclease Asp718 and 26 nucleotides complementary to the 3' non-translated sequence of the HDGF-2 gene. The amplified sequences are isolated from a 1% agarose gel using a commercially available kit ("Geneclean," BIO 101 Inc., La Jolla, Ca.). The fragment is then digested with the endonucleases BamHI and Asp718 and then purified again on a 1% agarose gel. This fragment is designated F2.

The vector pA2 (modification of pVL941 vector, discussed below) is used for the expression of the HDGF-2 protein using the baculovirus expression system (for review see: Summers, M.D. and Smith, G.E. 1987, A manual of methods for baculovirus vectors and insect cell culture procedures, Texas Agricultural Experimental Station Bulletin NO:1555). This expression vector contains the strong polyhedrin promoter of the Autographa californica nuclear polyhedrosis virus (AcMNPV) followed by the recognition sites for the restriction endonucleases . The polyadenylation site of the simian virus (SV)40 is used for efficient polyadenylation. For an easy selection of recombinant viruses the beta-galactosidase gene from E.coli is inserted in the same orientation as the polyhedrin promoter followed by the polyadenylation signal of the polyhedrin gene. The polyhedrin sequences are flanked at both sides by viral sequences for the cell-mediated homologous recombination of cotransfected wild-type viral DNA. Many other baculovirus

vectors could be used in place of pA2 such as pRG1, pAc373, pVL941 and pAcIM1 (Luckow, W.A. and Summers, M.D., Virology, 170:31-39).

The plasmid is digested with the restriction enzymes and then dephosphorylated using calf intestinal phosphatase by procedures known in the art. The DNA is then isolated from a 1% agarose gel using the commercially available kit ("Geneclean" BIO 101 Inc., La Jolla, Ca.). This vector DNA is designated V2.

Fragment F2 and the dephosphorylated plasmid V2 are ligated with T4 DNA ligase. E.coli XL-1 Blue cells are then transformed and bacteria identified that contained the plasmid (pBac-HDGF-2) with the HDGF-2 gene using the enzymes BamHI and Asp718. The sequence of the cloned fragment is confirmed by DNA sequencing.

5  $\mu$ g of the plasmid pBac-HDGF-2 is cotransfected with 1.0  $\mu$ g of a commercially available linearized baculovirus ("BaculoGold™ baculovirus DNA", Pharmingen, San Diego, CA.) using the lipofection method (Felgner et al. Proc. Natl. Acad. Sci. USA, 84:7413-7417 (1987)).

1 $\mu$ g of BaculoGold™ virus DNA and 5  $\mu$ g of the plasmid pBac-HDGF-2 are mixed in a sterile well of a microtiter plate containing 50  $\mu$ l of serum free Grace's medium (Life Technologies Inc., Gaithersburg, MD). Afterwards 10  $\mu$ l Lipofectin plus 90  $\mu$ l Grace's medium are added, mixed and incubated for 15 minutes at room temperature. Then the transfection mixture is added dropwise to the Sf9 insect cells (ATCC CRL 1711) seeded in a 35 mm tissue culture plate with 1ml Grace's medium without serum. The plate is rocked back and forth to mix the newly added solution. The plate is then incubated for 5 hours at 27°C. After 5 hours the transfection solution is removed from the plate and 1 ml of Grace's insect medium supplemented with 10% fetal calf serum is added. The plate is put back into an incubator and cultivation continued at 27°C for four days.

After four days the supernatant is collected and a plaque assay performed similar as described by Summers and Smith (supra). As a modification an agarose gel with "Blue Gal" (Life Technologies Inc., Gaithersburg) is used which allows an easy isolation of blue stained plaques. (A detailed description of a "plaque assay" can also be found in the user's guide for insect cell culture and baculovirology distributed by Life Technologies Inc., Gaithersburg, page 9-10).

Four days after the serial dilution, the viruses are added to the cells and blue stained plaques are picked with the tip of an Eppendorf pipette. The agar containing the recombinant viruses is then resuspended in an Eppendorf tube containing 200  $\mu$ l of Grace's medium. The agar is removed by a brief centrifugation and the supernatant containing the recombinant baculovirus is used to infect Sf9 cells seeded in 35 mm dishes. Four days later the supernatants of these culture dishes are harvested and then stored at 4°C.

Sf9 cells are grown in Grace's medium supplemented with 10% heat-inactivated FBS. The cells are infected with the recombinant baculovirus V-HDGF-2 at a multiplicity of infection (MOI) of 2. Six hours later the medium is removed and replaced with SF900 II medium minus methionine and cysteine (Life Technologies Inc., Gaithersburg). 42 hours later 5  $\mu$ Ci of  $^{35}$ S-methionine and 5  $\mu$ Ci  $^{35}$ S cysteine (Amersham) are added. The cells are further incubated for 16 hours before they are harvested by centrifugation and the labelled proteins visualized by SDS-PAGE and autoradiography.

### Example 3

#### Expression of Recombinant HDGF-2 in COS cells

The expression of plasmid, CMV-HDGF-2 HA is derived from a vector pcDNA1/Amp (Invitrogen) containing: 1) SV40 origin of replication, 2) ampicillin resistance gene, 3) E.coli replication origin, 4) CMV promoter followed by a polylinker

region, a SV40 intron and polyadenylation site. A DNA fragment encoding the entire HDGF-2 precursor and a HA tag fused in frame to its 3' end is cloned into the polylinker region of the vector, therefore, the recombinant protein expression is directed under the CMV promoter. The HA tag correspond to an epitope derived from the influenza hemagglutinin protein as previously described (I. Wilson, et al., Cell, 37:767 (1984)). The infusion of HA tag to the target protein allows easy detection of the recombinant protein with an antibody that recognizes the HA epitope.

The plasmid construction strategy is described as follow:

The DNA sequence encoding HDGF-2, ATCC # \_\_\_\_, contained in the plasmid vector pBluescript was amplified by PCR with a pBluescript vector primer (T3) at the 5' end and a HDGF-2 specific primer at the 3' end of the HDGF-2 coding sequence containing an XhoI restriction site. After amplification via PCR, the resultant PCR product is digested with BamHI and XhoI and ligated into a modified pCDNA-1 vector containing the HA tag in frame following the XhoI restriction site. The resultant plasmid contains the 5' untranslated region of HDGF-2 followed by the entire coding sequence fused in frame to the HA tag at the C-terminus.

The ligation mixture is transformed into E. coli strain SURE (Stratagene Cloning Systems, La Jolla, CA) the transformed culture is plated on ampicillin media plates and resistant colonies are selected. Plasmid DNA is isolated from transformants and examined by restriction analysis for the presence of the correct fragment. For expression of the recombinant HDGF-2, COS cells are transfected with the expression vector by DEAE-DEXTRAN method (J. Sambrook, E. Fritsch, T. Maniatis, Molecular Cloning: A Laboratory Manual, Cold Spring Laboratory Press, (1989)). The expression of the HDGF-2-HA protein is detected by radiolabelling and immunoprecipitation method (E. Harlow, D. Lane, Antibodies:

A Laboratory Manual, Cold Spring Harbor Laboratory Press, (1988)). Cells are labelled for 8 hours with  $^{35}\text{S}$ -cysteine two days post transfection. Culture media are then collected and cells are lysed with detergent (RIPA buffer (150 mM NaCl, 1% NP-40, 0.1% SDS, 1% NP-40, 0.5% DOC, 50mM Tris, pH 7.5) (Wilson, I. et al., Id. 37:767 (1984))). Both cell lysate and culture media are precipitated with a HA specific monoclonal antibody. Proteins precipitated are analyzed on 15% SDS-PAGE gels.

#### Example 4

##### Expression via Gene Therapy

Fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in tissue-culture medium and separated into small pieces. Small chunks of the tissue are placed on a wet surface of a tissue culture flask, approximately ten pieces are placed in each flask. The flask is turned upside down, closed tight and left at room temperature over night. After 24 hours at room temperature, the flask is inverted and the chunks of tissue remain fixed to the bottom of the flask and fresh media (e.g., Ham's F12 media, with 10% FBS, penicillin and streptomycin, is added. This is then incubated at 37°C for approximately one week. At this time, fresh media is added and subsequently changed every several days. After an additional two weeks in culture, a monolayer of fibroblasts emerge. The monolayer is trypsinized and scaled into larger flasks.

pMV-7 (Kirschmeier, P.T. et al, DNA, 7:219-25 (1988) flanked by the long terminal repeats of the Moloney murine sarcoma virus, is digested with EcoRI and HindIII and subsequently treated with calf intestinal phosphatase. The linear vector is fractionated on agarose gel and purified, using glass beads.

The cDNA encoding a polypeptide of the present invention is amplified using PCR primers which correspond to the 5' and



3' end sequences respectively. The 5' primer containing an EcoRI site and the 3' primer further includes a HindIII site. Equal quantities of the Moloney murine sarcoma virus linear backbone and the amplified EcoRI and HindIII fragment are added together, in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The ligation mixture is used to transform bacteria HB101, which are then plated onto agar-containing kanamycin for the purpose of confirming that the vector had the gene of interest properly inserted.

The amphotropic pA317 or GP+am12 packaging cells are grown in tissue culture to confluent density in Dulbecco's Modified Eagles Medium (DMEM) with 10% calf serum (CS), penicillin and streptomycin. The MSV vector containing the gene is then added to the media and the packaging cells are transduced with the vector. The packaging cells now produce infectious viral particles containing the gene (the packaging cells are now referred to as producer cells).

Fresh media is added to the transduced producer cells, and subsequently, the media is harvested from a 10 cm plate of confluent producer cells. The spent media, containing the infectious viral particles, is filtered through a millipore filter to remove detached producer cells and this media is then used to infect fibroblast cells. Media is removed from a sub-confluent plate of fibroblasts and quickly replaced with the media from the producer cells. This media is removed and replaced with fresh media. If the titer of virus is high, then virtually all fibroblasts will be infected and no selection is required. If the titer is very low, then it is necessary to use a retroviral vector that has a selectable marker, such as neo or his.

The engineered fibroblasts are then injected into the host, either alone or after having been grown to confluence on cytodex 3 microcarrier beads. The fibroblasts now produce the protein product.

Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, within the scope of the appended claims, the invention may be practiced otherwise than as particularly described.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: Kunsch, ET AL.
- (ii) TITLE OF INVENTION: Human Hepatoma-Derived Growth Factor
- (iii) NUMBER OF SEQUENCES: 8
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: CARELLA, BYRNE, BAIN, GILFILLAN, CECCHI, STEWART & OLSTEIN
  - (B) STREET: 6 BECKER FARM ROAD
  - (C) CITY: ROSELAND
  - (D) STATE: NEW JERSEY
  - (E) COUNTRY: USA
  - (F) ZIP: 07068
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: 3.5 INCH DISKETTE
  - (B) COMPUTER: IBM PS/2
  - (C) OPERATING SYSTEM: MS-DOS
  - (D) SOFTWARE: WORD PERFECT 5.1
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER:
  - (B) FILING DATE: Filed Herewith
  - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: None
  - (B) FILING DATE: None
- (viii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: FERRARO, GREGORY D.
  - (B) REGISTRATION NUMBER: 36,134
  - (C) REFERENCE/DOCKET NUMBER: 325800-
- (x) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: 201-994-1700
  - (B) TELEFAX: 201-994-1744

## (2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS
  - (A) LENGTH: BASE PAIRS
  - (B) TYPE: NUCLEIC ACID
  - (C) STRANDEDNESS: SINGLE
  - (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: AMINO ACIDS

(B) TYPE: AMINO ACID

(C) STRANDEDNESS:

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PEPTIDE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

WHAT IS CLAIMED IS:

1. An isolated polynucleotide comprising a member selected from the group consisting of:
  - (a) a polynucleotide encoding the polypeptide as set forth in Figure 1;
  - (b) a polynucleotide which encodes a mature polypeptide having the amino acid sequence expressed by the DNA contained in ATCC Deposit No. \_\_\_\_\_;
  - (c) a polynucleotide capable of hybridizing to and which is at least 70% identical to the polynucleotide of (a) or (b); and
  - (d) a polynucleotide fragment of the polynucleotide of (a) or (b), or (c).
2. The polynucleotide of Claim 2 which encodes the polypeptide comprising amino acid 1 to 249 of SEQ ID NO:2.
3. A vector containing the polynucleotide of Claim 1.
4. A host cell genetically engineered with the vector of Claim 3.
5. A process for producing a polypeptide comprising: expressing from the host cell of Claim 4 the polypeptide encoded by said polypeptide.
6. A process for producing cells capable of expressing a polypeptide comprising genetically engineering cells with the vector of Claim 3.
7. A polypeptide selected from the group consisting of (i) a polypeptide having the deduced amino acid sequence of SEQ ID NO:2 and fragments, analogs and derivatives thereof; and (ii) a polypeptide encoded by the cDNA of ATCC Deposit No. \_\_\_\_\_ and fragments, analogs and derivatives of said polypeptide.

8. The polypeptide of Claim 13 wherein the polypeptide comprises the amino acids of Figure 1.
9. An antibody against the polypeptide of Claim 7.
10. A compound which inhibits activation of the polypeptide of claim 7.
11. A method for the treatment of a patient having need of HDGF-2 comprising: administering to the patient a therapeutically effective amount of the polypeptide of claim 7.
12. The method of Claim 11 wherein said therapeutically effective amount of the polypeptide is administered by providing to the patient DNA encoding said polypeptide and expressing said polypeptide in vivo.
13. A method for the treatment of a patient having need to inhibit a HDGF-2 polypeptide comprising: administering to the patient a therapeutically effective amount of the compound of Claim 10.
14. A process for diagnosing in a patient a disease or a susceptibility to a disease related to an under-expression of the polypeptide of claim 7 comprising:  
determining a mutation in a nucleic acid sequence encoding said polypeptide in a sample derived from a patient.
15. A diagnostic process comprising:  
analyzing for the presence of the polypeptide of claim 7 in a sample derived from a host.
16. A method for identifying compounds which is an agonist of the polypeptide of claim 7 comprising:

contacting a cell expressing on the surface thereof a receptor for the polypeptide, said receptor being associated with a second component capable of providing a detectable signal in response to the binding of a compound to said receptor, with a compound under conditions to permit binding to the receptor; and

determining whether the compound binds to and activates the receptor by detecting the presence of a signal generated from the interaction of the compound with the receptor.

17. A method for identifying compounds which bind to and inhibit activation of the polypeptide of claim 13 comprising:

contacting a cell expressing on the surface thereof an HDGF-2 receptor polypeptide, said receptor being associated with a second component capable of providing a detectable signal in response to the binding of a compound to said receptor polypeptide, with HDGF-2 polypeptide and a compound to be screened under conditions to permit binding to the receptor polypeptide; and

determining whether the compound inhibits the HDGF-2 polypeptide by detecting the absence of a signal.

1 / 4

## FIG. 1A

1 GAATTCGTCTCTTAGGGTGGTTGGGTGTAAGATGGCGGCTGTGAGTCTGCGGCTCGGC  
M A A V S L R L G

61 GACTTGGTGTGGGGAACCTCGGCCGATATCCTCCTTGGCCAGGAAGATTGTTAATCCA  
D L V W G K L G R Y P P W P G K I V N P

121 CCAAGGACTTGAAGAAACCTCGCGGAAGAAATGCTTCTTTGTGAAATTTTGTGAACA  
P K D L K K P R G K K C F F V K F F G T

181 GAAGATCATGCCCTGGATCAAAGTGAACAGCTGAAGCCATATCATGCTCATAAAGAGGAA  
E D H A W I K V E Q L K P Y H A H K E E

241 ATGATAAAAATTAAAGGGTAAACGATTCCAGCAAGCGGTAGATGCTGTCGAAGAGTTC  
M I K I N K G K R F Q Q A V D A V E E F

301 CTCAGGAGAGCCAAAGGGAAGACCAGACGTCAATCCACAAATTTCTGTGACAAAGAAT  
L R R A K G K D Q T S S H N S S D D K N

361 CGACGTAATTCCAGTGAGGAGAGAAAGTAGGCCAAACTCAGGTGATGAGAAGCGCAAACCTT  
R R N S S E E R S R P N S G D E K R K L

421 AGCCTGTCTGAAGGGAAGGTGAAGAAGAACATGGGAGAAGGAAGAGGGTGTCTTCA  
S L S E G K V K K N M G E G K K R V S S

481 GGCTCTTCAGAGAGAGGCTCCAAATCCCCCTCTGAAAAGAGCCCCAAGAGCAAAGTCCCCGG  
G S S E R G S K S P L K R A Q E Q S P R

541 AAGCGGGTCTGGCCCCCAAGGATGAGAAGGATCTCACCATCCCCGGAGTCTAGTACCGTG  
MATCH WITH FIG. 1B



2 / 4

## FIG. 1B

MATCH WITH FIG. 1A

K R G R P P K D E K D L T I P E S S T V

601 AAGGGATGATGGCCGACCGATGGCCCGGTTAAATGGCAGCCAAACGCAAGCAGCCT  
 K G M M A G P M A A F K W Q P T A S E P

661 GTTAAAGATGCAGATCCTCATTTCCATCATTTCTGCTAAGCCAAACAGAGAAGCCAGCT  
 V K D A D P H F H F L L S Q T E K P A

721 GTCTGTTACCAGGCAATCACGAAGAAGTTGAAAATATGTGAAGACCTTCCTTCTCCTAGG  
 V C Y Q A I T K K L K I C E D L L L P R

781 TGAAGTGGCAATGCAGCCAGATGATGCTGATCGTGAACATGGTCCAAGGGAGCTTCAT  
 841 GGCCACTATTGCCAGGGGCTGACCTGGCCCCAGGTGACAGGCCAGTCCCAGCAGACACT  
 901 CTTGGACATCCTCAATCAGGGACAGTTGGCCAGCATCTTCTGGACCAGAAAGTGCCAAA  
 961 TATCCTGCAAGGAACTTTAAGCCTGATTTCTACCTGAAATACATTCAGAAAGGATCTCCG  
 1021 CTTAGCCATTGCCGTGGTGATGCGGTCAACCATCCGACTCCCATGGCAGCTGCAGCAAA  
 1081 TGAGGTGTACAAAAGAGCCAGGCGCTGGACCAAGTGTGACAAACGATATGTCCGCCGTGA  
 1141 CCGAGCCTACATACACTAAGCTGTGACACACCCGCCCTCACCCTCCCAATCCCCCTCTG  
 1201 ACCCCCTCTTCCTCACATGGGGTCGGGGCTGGGAGTTTCATTCTGGTACCAGCCCCACCT  
 1261 ATCTCCATTTCCTTTTATACAGACTTTTGAGACTTGCCATCAGCACACACAGCAGCAC  
 1321 CCTTCCCCCTGAGTCGGTGGGAGGGGACAAAGTGTGAGCAGGATTTGGCGTGTGGAAAGC  
 1381 TCTTGAGCTGGGCACCTGGCCCCCGGACGAGGTGGTGTGTGTTTCACACACACACACA  
 1441 CACACACACACACACACACAGGCTCTCGCCCCCAGGATAGAAAGCTGCCCAGAAACTG  
 1501 CTGCCCTGGCTTTTCTTCCGAGCTTGCTTATCTCAAACCCCTTCCAGTCAAGGAACCT  
 1561 AGAATCAGCAACGAGAGTTGGAAGCCTTCCCACAGCTTCCCCCAGAGCGAAGGCTGTA  
 1621 GTCATGTCCCATCCCCACTGGATTCCCTACAAAGGAGAGGCCCTTGGGCCCCAGATGAGCC  
 1681 AGTACAGACTCCAGACAGAGGGGCCCTTGGGGCCCTCCAACTCAGGTGATGAGCTGAGA  
 1741 AAGATGTTACGTCCTAAGCGTCCAGTGTGACCCAGCGCTCCATAGACGCCCTTTGTGAAC

MATCH WITH FIG. 1C

## FIG. 1C

MATCH WITH FIG. 1B

1801 TGAAGAAGAGACTGGCAGAGTCCCGAGAAGATGGGGCCCTGGCTTTCCAGGGAGTGCAGCA  
1861 AGCAGCCGGCCTGCAGGTGAGCATGGAGGCCCGGCCCTCACCGCCCTCGAAGCCATGCCCC  
1921 AGATGCCACTGCCACAGCGGGCGCTCGCTCCCTAGGCTGTTTATAGTATTTGGATTG  
1981 CATTCATCCCTTGGGAGGGAGTCCCTCAGGGCCACTAGTGTAGAGCCAAAGAGGAGTGGGG  
2041 GTTGGGGCGCTCCTTTCTGTTTCCGTTAGGCCACAGACTCTTCACCTGGCTCTGACTTA  
2101 CCTCGGTCCCTCCAGTGGTCCACCTTCTCCACCTGCCCTGCCAAGTCCCTGCATG  
2161 CCCACCGCTCTCCATCCTCCCTCTCCCTCTCCCTCCCGTGGAGACAGTATTTCTTTC  
2221 TGTCTGTCCCTTTGGCCCCAGACCCAGCCCTGACCAACGATGAGCATTTCTTAGGCTCAGCT  
2281 CTTGATACGGAAACGAGTGTCTTCACTCCAGCCAGCATCATGCTCTTCGGTGTCTCCCGG  
2341 GCCCGGGTCTGTCTGGGAGGGAAGAGAACTGGGCCCTGACCTACCTGAAC TGACTGGCCCT  
2401 CCGAGGTGGTCTGGGACATCCTAGAGGCCCTACATTTGTCTCTGGATAGGGACCGGGG  
2461 GGGCTTGGAAATGTTSCAAAAAAGTTACCCAGGATGTCAGTTTTATATCCCTCT  
2521 GCATGGGTTGGATTTTCCAAAAATCATAAATTTGCAGAAAGGAGCCAGCATTTACGATGCA  
2581 ATATGTAATTATATATAGGGTGGCCACACTAGGGGGGTCTTCCCCCTCACAGCTTT  
2641 GGCCCTTTTCAGAGATTAGAAACTGGGTTAGAGGATTGCAGAAAGACGAGTGGGGGAGGG  
2701 CAGGGAAGATGCCCTGTCTGGGTTTATAGCACAGTTTCATTTACATGGGATTTTGAAGCATTT  
2761 CTGTCTGAACACAAAGCCGTGTTCTAGTCCCTGGCGGAACACACTGGGGTGGGGCGGGGG  
2821 AAGATGCGGTAATGAACCCGGTTAGTCAATTTGTCTTAATAATGTTGACAAATCTGTAA  
2881 AGTTCCCTTTTATGAATAATTTCTGTTTAAGCTATTTTCACCTTTCTTTTGAAATCTCTCC  
2941 TTTTAAGGAGAAATAATGTGACACTTTGTGAAAAAGCTTGTGAAGAAAGCCCTTCTTTT  
3001 CTTTAAACCTTTAAATGACAAATCTAGGTAATTAAGTTGTGAATTTTATTTTGTCTTT  
3061 GTTTTAAATGAACATTTGTCTTTTCAGAAATAGGATTTGTGTGATAATGTTTAAATGGSAAA  
3121 ACAAAACATGATTTTGTGCAATTAACAAAGCTACTGCAAGGAAAAATAAAACACTTCTTGG  
3181 TAACAAAAAATAAAAAA 3202



# INTERNATIONAL SEARCH REPORT

Application No.  
PCT/US95/06731

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12N 1/21, 5/10, 15/09, 15/12, 15/63, 15/70; C07K 14/475

US CL : 435/69.1, 320.1, 252.3, 240.2; 514/2; 530/350; 536/23.5

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/69.1, 320.1, 252.3, 240.2; 514/2; 530/350; 536/23.5

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Journal Biol. Chem. Volume 269, Number 40, issued 07 October 1994, Nakamura et al., "Molecular Cloning of Complementary DNA for a Novel Human Hepatoma-derived Growth Factor," pages 25143-25149, especially Figure 2, page 25146.	1-6
A	Clin. Chim. Acta, Volume 183, issued 1989, Nakamura et al., "Partial Purification and Characterization of Human Hepatoma-derived Growth Factor," pages 273-284, especially pages 277-279.	7-8
X, O	EMBL-Genbank, issued 17 April 1995, Hillier et al., Locus R19745, Definition yg40c08.r1 Homo sapiens cDNA clone 34940 5', see entire sequence.	1, 3-6

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A document defining the general state of the art which is not considered to be of particular relevance	*X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E earlier document published on or after the international filing date	*Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z document member of the same patent family
*O document referring to an oral disclosure, use, exhibition or other means	
*P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

26 SEPTEMBER 1995

Date of mailing of the international search report

13 OCT 1995

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Authorized officer

KAREN E. BROWN aco

Facsimile No. (703) 305-3230

Telephone No. (703) 308-0196

# INTERNATIONAL SEARCH REPORT

Application No.  
PCT/US93/06731

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EMBL-Genbank, issued 30 July 1993, Genexpress, Locus HSBA6E042, Description H. sapiens partial cDNA sequence, clone A6E04, see entire sequence.	1, 3-6
X	EMBL, Genbank, issued 15 February 1995, Genexpress, Locus HSCOZA111, Description H. sapiens partial cDNA sequence, clone c-Oza11, see entire sequence.	1, 3-6
X	EMBL-Genbank, issued 08 February 1995, Hillier et al., Locus T47923, Definition yb18c09.s1 Homo sapiens cDNA clone 71536 3', see entire sequence.	1, 3-6
X	EMBL-Genbank, issued 13 May 1995, Hillier et al., Locus HS164111, Description yg40c08.s1 Homo sapiens cDNA clone 34940 3', see entire document sequence.	1, 3-6
X	Nature Genetics, Volume 4, issued July 1993, Adams et al., "3,400 New Expressed Sequence Tags Identify Diversity of Transcripts in Human Brain," pages 256-267, see Table 2.	1, 3-6
X	EMBL-Genbank, published 30 June 1993, Adams et al., Locus T05906, Description Homo sapiens cDNA clone HFBD141, see entire sequence.	1, 3-6

# INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/US93/06731

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
  
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
  
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:  
1-8

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

ii. application No.  
PCT/US95/06731

## BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

Group I, claims 1-8, drawn to polynucleotides, vectors, host cells, process of producing polypeptides, and polypeptides for HDGF-2.

Group II, claim 9, drawn to antibody against HDGF-2.

Group III, claims 10 and 13, drawn to an inhibitor of activation of HDGF-2 and a method of using said inhibitor.

Group IV, claims 11-12, drawn to a method of treating patients with HDGF-2 using DNA or protein.

Group V, claim 14, drawn to a method of diagnosing a disease or susceptibility to a disease by determining a mutation in the HDGF-2 DNA.

Group VI, claim 15, drawn to a method of determining the presence of the HDGF-2 protein.

Group VII, claims 16-17, drawn to methods of identifying agonists or antagonists to HDGF-2. The inventions listed as Groups I-VII do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

None of the polypeptides and polynucleotides of Group I, the antibody of Group II, or the inhibitor of Group II share structural or functional features, and each is defined by special technical structural and functional features which distinguishes each from the others.

None of the methods of treating patients of Group IV, the method of diagnosing disease of Group V, the method of determining polypeptide levels of Group VI or the method of identifying agonists or antagonists of Group VII share a special technical feature with any other group because each method is distinguished from each of the others by its unique process steps.

None of the products of any of Groups I-III are used in or produced by the methods of any of Groups V-VII.

The methods of Group I do not share any special technical features with the methods of Group IV because their different process steps are technical features which distinguish them.

Accordingly, the claims are not linked by a special technical feature within the meaning of PCT Rule 13.2 so as to form a single inventive concept.